




#### **Extreme Rainfall in Zhengzhou**

From 18:00 on July 18<sup>th</sup> to 0:00 on July 21<sup>st</sup>, 2021, a rare and sustained heavy precipitation occurred in **Zhengzhou**, Henan province. In three days, an entire **year's-worth** of rain fell on this city.

From 16:00 to 17:00 on July 20th, 2021, the rainfall in Zhengzhou reached **201.9 mm in one hour**.





More than 100 West Lakes poured into Zhengzhou in one hour



#### **Related News**

#### **BBC**

# China floods: Thousands evacuated as rail stations and roads submerged

1 hour ago | China

#### Guardian

Heavy flooding hits central China, affecting tens of millions

#### **New York Times**


ASIA PACIFIC

Floods in China Leave Many Stranded

#### **Extreme Rainfall in Zhengzhou**



Here are **some photos** pictured by one of my friends when he was stuck in Zhengzhou.



A: View from Zhengzhou East High-speed Railway Station



B: Life supplies delivered to Zhengzhou



C: Volunteers played a very important role



#### **Experience in Beijing**





 $PM_{10}$  concentrations in most areas exceeded 2000  $\mu G/m3$ 

# A sandstorm that hit Beijing in March pushed air pollution **levels off the charts**, turned the sky yellow and reduced the sun to a blue dot.

Thick dust carrying extremely high levels of hazardous particles blows in from drought-hit **Mongolia**.




Fig 2: a natural-color image shows trajectory of dust (NASA)

# **SANDSTORM**



A photo on March 15<sup>th</sup> 2021, on PKU campus

"It's of the greatest intensity over 10 years!"

# **Experience in Beijing**





PKU campus shrouded in the snow

## Extreme cold

2021.1.7: -19.6 °C, the second lowest winter temperature since 1951


2021.10.17: -0.2 °C ,the lowest since 1969. The first temperature below zero was 20 days earlier.

2021.11.6: first snow, which is much earlier than last year (11.21)

# **Extreme rainfall**

Long Flood season 122 DAYSStrong


- **1.** 10 times over **25 mm/d**, 2 times over **50 mm/d**
- 2. 792mm, 90% higher than annual data, 70% higher than the data in recent 10 years
- 3. 400.4mm in July, the most since monitoring began (1951)



# Changes in heavy precipitation in the world



(b) Synthesis of assessment of observed change in **heavy precipitation** and confidence in human contribution to the observed changes in the world's regions



Source: The Sixth Assessment Report of the Intergovernmental Panel on Climate Change

Type of observed change in heavy precipitation

Increase (19)

Decrease (0)

Low agreement in the type of change (8)

Limited data and/or literature (18)


Confidence in human contribution to the observed change

- • High
- Medium
- Low due to limited agreement
- Low due to limited evidence

#### Heavy precipitation over land

#### 10-year event

Frequency and increase in intensity of heavy 1-day precipitation event that occurred **once in 10 years** on average **in a climate without human influence** 






# Heavy precipitation (frequency and intensity)

- Projected changes are shown at global warming levels of 1°C, 1.5°C, 2°C, and 4°C and are relative to 1850–1900,9 representing a climate without human influence. The figure depicts frequencies and increases in intensity of 10- or 50-year extreme events from the base period (1850–1900) under different global warming levels.
- Extreme precipitation events are defined as the daily precipitation amount over land that was exceeded on average once in a decade during the 1850–1900 reference period.

Source: The Sixth Assessment Report of the Intergovernmental Panel on Climate Change

#### **Economic Issue**





- ec
  - Transformation risks associated with revaluation of carbon intensive assets

The blue bar here means the loss annually and the red line means the average loss over the last 30 years (Units: One hundred million yuan)

#### Features of extreme weather:

- Wide variety
- high frequency
- long duration
- strong seasonality
- obvious regionality


- Extreme weather
  - Bring greater economic costs
- Financial system
  - Physical risks associated with economic losses caused by climate related events

- Agricultural sector
- probably one of the most sensitive sectors to climate change In 2020, the direct economic loss reached 368.1 billion yuan



#### **Economic Issue**





IEA, 2021.

Notes: Bubble area represents total energy-related and process-related CO<sub>2</sub> emissions.


CO<sub>2</sub> emissions intensity of primary energy demand relative to CO<sub>2</sub> emissions per capita by country/region, 2000 and 2020

 Despite impressive growth in renewables since 2000, China remains heavily dependent on fossil fuels, with coal alone still meeting 60% of its total primary energy needs.



### **Agriculture Issue**







#### **Agriculture Issue**



Many diseases and pests began to evolve a unique hibernation mechanism, and the survival time gradually increased plant diseases & insect pests Many of the northern diseases and insects were brought to the south without corresponding natural enemies to contain them Large-scale precipitation will cause landslides and mountain floods, resulting in the breach of the reservoir dike extreme weather Climate Change Extreme droughts can reduce crop yields Example: Since the 1990s, the climate in northeast China has changed significantly, and rice can no longer be light resources; Planting area and system. grown in some areas, while the planting range of warm-loving plants in northwest China has expanded.

## **Agriculture Issue**




China is making great efforts in agricultural production to mitigate the impact of global climate change on food production.

Academician Yuan Longping (1930.9-2021.5)
"Father of Hybrid Rice"

"I have two dreams:
The dream of enjoying the cool under the grain.
The dream of covering the world with hybrid rice."

For People in China, for People of the World.




On October 26, 2021, nanfan public experimental base in Sanya city, Hainan province announced the results of double season rice yield measurement: Hainan has achieved the goal of 1500 kg per unit area of double season rice, creating the highest local double season rice yield per mu record!


#### **Health Issue**



In 49 large cities in the United States, changes in extreme hot and extreme cold temperatures are projected to result in more than 9,000 additional premature deaths per year under a higher scenario by the end of the century.



Mean summer warming relative to the 1986–2005 average




Change in the number of heatwave exposure compared with the historical average number of events (1986–2005 average)



#### **Health Issue**



#### Ten Climate Hazards And six aspects of human systems







Fig. 1. Climate change, human infectious diseases, and human society.



# **Adaptation**



·Some countries have taken national health climate adaptation strategies or plans.





#### Adaptation

#### Local efforts including:

- Altering urban design (for example, by using cool roofs, tree shades, and green walkways)
- Improving water management (for example, via desalination plants or watershed protection)

can provide health and social justice benefits, elicit neighborhood participation, and increase resilience for specific populations, such as outdoor workers.

#### Key Message 3



#### Adaptation Reduces Risks and Improves Health

Proactive adaptation policies and programs reduce the risks and impacts from climate-sensitive health outcomes and from disruptions in healthcare services. Additional benefits to health arise from explicitly accounting for climate change risks in infrastructure planning and urban design.





# Thanks!



